

SOME REFINEMENTS OF EXISTENCE RESULTS FOR SPDES DRIVEN BY WIENER PROCESSES AND POISSON RANDOM MEASURES

STEFAN TAPPE

ABSTRACT. We provide existence and uniqueness of global (and local) mild solutions for a general class of semilinear stochastic partial differential equations driven by Wiener processes and Poisson random measures under local Lipschitz and linear growth (or local boundedness, resp.) conditions. The so-called “method of the moving frame” allows us to reduce the SPDE problems to SDE problems.

1. INTRODUCTION

Semilinear stochastic partial differential equations (SPDEs) on Hilbert spaces, being of the type

$$(1.1) \quad \begin{cases} dZ_t &= (AZ_t + \alpha(t, Z_t))dt + \sigma(t, Z_t)dW_t \\ Z_0 &= z_0, \end{cases}$$

have widely been studied in the literature, see e.g. [4, 23, 27, 10]. In equation (1.1), A denotes the generator of a strongly continuous semigroup, and W is a trace class Wiener process. In view of applications, this framework has been extended by adding jumps to the SPDE (1.1). More precisely, consider a SPDE of the type

$$(1.2) \quad \begin{cases} dZ_t &= (AZ_t + \alpha(t, Z_t))dt + \sigma(t, Z_t)dW_t \\ &+ \int_E \gamma(t, Z_{t-}, x)(\mu(dt, dx) - F(dx)dt) \\ Z_0 &= z_0, \end{cases}$$

where μ denotes a Poisson random measure on some mark space (E, \mathcal{E}) with $dt \otimes F(dx)$ being its compensator. SPDEs of this type have been investigated in [20, 7], see also [17, 18, 16, 21, 1, 22], where SPDEs with jump noises have been studied.

The goal of the present paper is to extend results and methods for SPDEs of the type (1.2) in the following directions:

- We consider more general SPDEs of the form

$$(1.3) \quad \begin{cases} dZ_t &= (AZ_t + \alpha(t, Z_t))dt + \sigma(t, Z_t)dW_t \\ &+ \int_B \gamma(t, Z_{t-}, x)(\mu(dt, dx) - F(dx)dt) \\ &+ \int_{B^c} \gamma(t, Z_{t-}, x)\mu(dt, dx) \\ Z_0 &= z_0, \end{cases}$$

where $B \in \mathcal{E}$ is a set with $F(B^c) < \infty$. Then, the integral \int_B represents the small jumps, and \int_{B^c} represents the large jumps of the solution process. Similar SDEs have been considered in finite dimension in [14, Sec. II.2.c] and in infinite dimension in [3].

Date: 6 August, 2012.

2010 Mathematics Subject Classification. 60H15, 60G57.

Key words and phrases. Semilinear SPDE with jumps, Poisson random measure, mild solution, method of the moving frame.

The author is grateful to an anonymous referee for valuable comments and suggestions.

- We will prove the following results (see Theorem 4.5) concerning existence and uniqueness of local and global mild solutions to (1.3):
 - (1) If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz and of linear growth, then existence and uniqueness of global mild solutions to (1.3) holds.
 - (2) If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz and locally bounded, then existence and uniqueness of local mild solutions to (1.3) holds.
 - (3) If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz, then uniqueness of mild solutions to (1.3) holds.

In particular, the result that local Lipschitz and linear growth conditions ensure existence and uniqueness of global mild solutions does not seem to be well-known for SPDEs, as most of the mentioned references impose global Lipschitz conditions. An exception is the reference [27], where the author treats Wiener process driven SPDEs of the type (1.1), even on 2-smooth Banach spaces, and provides existence and uniqueness under local Lipschitz and linear growth conditions. In [27], the crucial assumption on the operator A is that it generates an analytic semigroup, while our results hold true for every pseudo-contractive semigroup.

- We reduce the proofs of these SPDE results to the analysis of SDE problems. This is due to the “method of the moving frame”, which has been presented in [7]. As a direct consequence, we obtain that any mild solution to (1.3) is càdlàg.

As just mentioned, we shall utilize the “method of the moving frame” from [7], which allows us to reduce the SPDE problems to SDE problems. Therefore, we will be concerned with SDEs in Hilbert spaces being of the type

$$(1.4) \quad \begin{cases} dY_t &= a(t, Y_t)dt + b(t, Y_t)dW_t + \int_B c(t, Y_{t-}, x)(\mu(dt, dx) - F(dx)dt) \\ &\quad + \int_{B^c} c(t, Y_{t-}, x)\mu(dt, dx) \\ Y_0 &= y_0. \end{cases}$$

By using the technique of interlacing solutions at jump times (which, in particular cases has been applied e.g. in [2, Sec. 6.2] and [21, Sec. 9.7]), we can reduce the SDE (1.4) to SDEs of the form

$$(1.5) \quad \begin{cases} dY_t &= a(t, Y_t)dt + b(t, Y_t)dW_t + \int_B c(t, Y_{t-}, x)(\mu(dt, dx) - F(dx)dt) \\ Y_0 &= y_0 \end{cases}$$

without large jumps, and for those SDEs suitable techniques and results are available in the literature. This allows us to derive existence and uniqueness results for the SDE (1.4), which are subject to the regularity conditions described above. We point out that the reference [3] also studies Hilbert space valued SDEs of the type (1.4) and provides an existence and uniqueness result considerably going beyond the classical results which impose global Lipschitz conditions. In Section 3.3, we provide a comparison of our existence and uniqueness result for SDEs of the type (1.4) with that from [3].

The remainder of this paper is organized as follows: In Section 2 we provide the required preliminaries and notation. In Section 3 we prove existence and uniqueness results for (local) strong solutions to SDEs of the form (1.4), and in Section 4 we prove existence and uniqueness results for (local) mild solutions to SPDEs of the form (1.3) by using the “method of the moving frame”.

2. PRELIMINARIES AND NOTATION

In this section, we provide the required preliminary results and some basic notation.

Throughout this text, let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ with $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$ be a filtered probability space satisfying the usual conditions.

Let U be a separable Hilbert space and let $Q \in L(U)$ be a nuclear, self-adjoint, positive definite linear operator. Then, there exist an orthonormal basis $(e_j)_{j \in \mathbb{N}}$ of U and a sequence $(\lambda_j)_{j \in \mathbb{N}} \subset (0, \infty)$ with $\sum_{j \in \mathbb{N}} \lambda_j < \infty$ such that

$$Qe_j = \lambda_j e_j \quad \text{for all } j \in \mathbb{N},$$

namely, the λ_j are the eigenvalues of Q , and each e_j is an eigenvector corresponding to λ_j . The space $U_0 := Q^{1/2}(U)$, equipped with the inner product

$$\langle u, v \rangle_{U_0} := \langle Q^{-1/2}u, Q^{-1/2}v \rangle_U,$$

is another separable Hilbert space and $(\sqrt{\lambda_j}e_j)_{j \in \mathbb{N}}$ is an orthonormal basis. Let W be an U -valued Q -Wiener process, see [4, p. 86, 87]. For another separable Hilbert space H , we denote by $L_2^0(H) := L_2(U_0, H)$ the space of Hilbert-Schmidt operators from U_0 into H , which, endowed with the Hilbert-Schmidt norm

$$\|\Phi\|_{L_2^0(H)} := \left(\sum_{j \in \mathbb{N}} \|\Phi(\sqrt{\lambda_j}e_j)\|^2 \right)^{1/2}, \quad \Phi \in L_2^0(H)$$

itself is a separable Hilbert space.

Let (E, \mathcal{E}) be a measurable space which we assume to be a Blackwell space (see [6, 11]). We remark that every Polish space with its Borel σ -field is a Blackwell space. Furthermore, let μ be a time-homogeneous Poisson random measure on $\mathbb{R}_+ \times E$, see [14, Def. II.1.20]. Then its compensator is of the form $dt \otimes F(dx)$, where F is a σ -finite measure on (E, \mathcal{E}) .

For the following definitions, let τ be a finite stopping time.

- We define the new filtration $\mathbb{F}^{(\tau)} = (\mathcal{F}_t^{(\tau)})_{t \geq 0}$ by

$$\mathcal{F}_t^{(\tau)} := \mathcal{F}_{\tau+t}, \quad t \geq 0.$$

- We define the new U -valued process $W^{(\tau)}$ by

$$W_t^{(\tau)} := W_{\tau+t} - W_\tau, \quad t \geq 0.$$

- We define the new random measure $\mu^{(\tau)}$ on $\mathbb{R}_+ \times E$ by

$$\mu^{(\tau)}(\omega; B) := \mu(\omega; B_{\tau(\omega)}), \quad \omega \in \Omega \text{ and } B \in \mathcal{B}(\mathbb{R}_+) \otimes \mathcal{E},$$

where we use the notation

$$B_\tau := \{(t + \tau, x) \in \mathbb{R}_+ \times E : (t, x) \in B\}.$$

Then, $W^{(\tau)}$ is a $\mathbb{F}^{(\tau)}$ -adapted Q -Wiener process and $\mu^{(\tau)}$ is a time-homogeneous Poisson random measure relative to the filtration $\mathbb{F}^{(\tau)}$ with compensator $dt \otimes F(dx)$, cf. [8, Lemma 4.6].

2.1. Lemma. *Let ϱ be another stopping time. Then, the mapping $(\varrho - \tau)^+$ is a $\mathbb{F}^{(\tau)}$ -stopping time.*

Proof. For every $t \in \mathbb{R}_+$ we have

$$\{(\varrho - \tau)^+ \leq t\} = \{\varrho - \tau \leq t\} = \{\varrho \leq \tau + t\} \in \mathcal{F}_{\tau+t} = \mathcal{F}_t^{(\tau)},$$

showing that $(\varrho - \tau)^+$ is a $\mathbb{F}^{(\tau)}$ -stopping time. \square

Denoting by $\mathcal{P}^{(\tau)}$ the predictable σ -algebra relative to the filtration $\mathbb{F}^{(\tau)}$, we have the following auxiliary result.

2.2. Lemma. *The following statements are true:*

(1) *The mapping*

$$\theta_\tau : \Omega \times \mathbb{R}_+ \rightarrow \Omega \times \mathbb{R}_+, \quad \theta_\tau(\omega, t) := (\omega, \tau(\omega) + t)$$

is $\mathcal{P}^{(\tau)}$ - \mathcal{P} -measurable.

(2) *The mapping*

$$\vartheta_\tau : \Omega \rightarrow \Omega \times \mathbb{R}_+, \quad \vartheta_\tau(\omega) := (\omega, \tau(\omega))$$

is \mathcal{F}_τ - \mathcal{P} -measurable.

Proof. According to [14, Thm. I.2.2], the system of sets

$$\{A \times \{0\} : A \in \mathcal{F}_0\} \cup \{\llbracket 0, \varrho \rrbracket : \varrho \text{ is a stopping time}\}$$

is a generating system of the predictable σ -algebra \mathcal{P} . For any set $A \in \mathcal{F}_0$ we have

$$\theta_\tau^{-1}(A \times \{0\}) = (A \cap \{\tau = 0\}) \times \{0\} \in \mathcal{P}^{(\tau)}.$$

Furthermore, for any \mathbb{F} -stopping time ϱ we have

$$\begin{aligned} \theta_\tau^{-1}(\llbracket 0, \varrho \rrbracket) &= \theta_\tau^{-1}(\{(\omega, t) \in \Omega \times \mathbb{R}_+ : 0 \leq t \leq \varrho(\omega)\}) \\ &= \{(\omega, t) \in \Omega \times \mathbb{R}_+ : 0 \leq \tau(\omega) + t \leq \varrho(\omega)\} \\ &= \{(\omega, t) \in \Omega \times \mathbb{R}_+ : 0 \leq t \leq \varrho(\omega) - \tau(\omega)\} \\ &= \llbracket 0, \varrho - \tau \rrbracket = \llbracket 0, (\varrho - \tau)^+ \rrbracket \setminus (\{\tau > \varrho\} \times \{0\}) \in \mathcal{P}^{(\tau)}, \end{aligned}$$

where, in the last step, we have used Lemma 2.1. This proves the first statement.

According to [14, Thm. I.2.2], the system of sets

$$\{A \times \{0\} : A \in \mathcal{F}_0\} \cup \{A \times (s, t] : s < t \text{ and } A \in \mathcal{F}_s\}$$

is a generating system of the predictable σ -algebra \mathcal{P} . For any set $A \in \mathcal{F}_0$ we have

$$\vartheta_\tau^{-1}(A \times \{0\}) = A \cap \{\tau = 0\} \in \mathcal{F}_0 \subset \mathcal{F}_\tau.$$

Furthermore, for all $s, t \in \mathbb{R}_+$ with $s < t$ and $A \in \mathcal{F}_s$ we have

$$\vartheta_\tau^{-1}(A \times (s, t]) = A \cap \{s < \tau\} \cap \{\tau \leq t\} \in \mathcal{F}_\tau,$$

establishing the second statement. \square

Let us further investigate the Poisson random measure μ . According to [14, Prop. II.1.14], there exist a sequence $(\kappa_n)_{n \in \mathbb{N}}$ of finite stopping times with $\llbracket \kappa_n \rrbracket \cap \llbracket \kappa_m \rrbracket = \emptyset$ for $n \neq m$ and an E -valued optional process ξ such that for every optional process $\gamma : \Omega \times \mathbb{R}_+ \times E \rightarrow H$, where H denotes a separable Hilbert space, and all $0 \leq t \leq u$ with

$$\mathbb{P}\left(\int_t^u \int_E \|\gamma(s, x)\|^2 \mu(ds, dx) < \infty\right) = 1$$

we have

$$(2.1) \quad \int_t^u \int_E \gamma(s, x) \mu(ds, dx) = \sum_{n \in \mathbb{N}} \gamma(\kappa_n, \xi_{\kappa_n}) \mathbb{1}_{\{t < \kappa_n \leq u\}}.$$

Let $B \in \mathcal{E}$ be a set with $F(B^c) < \infty$. We define the mappings $\varrho_k : \Omega \rightarrow \overline{\mathbb{R}}_+$, $k \in \mathbb{N}_0$ as

$$\varrho_k := \inf\{t \geq 0 : \mu([0, t] \times B^c) = k\}, \quad k \in \mathbb{N}_0.$$

2.3. Lemma. *The following statements are true:*

- (1) *For each $k \in \mathbb{N}$ the mapping ϱ_k is a finite stopping time.*
- (2) *We have $\varrho_0 = 0$ and $\mathbb{P}(\varrho_k < \varrho_{k+1}) = 1$ for all $k \in \mathbb{N}_0$.*
- (3) *We have $\mathbb{P}(\varrho_k \rightarrow \infty) = 1$.*

Proof. This follows from [9, Lemma A.19]. \square

3. EXISTENCE AND UNIQUENESS OF STRONG SOLUTIONS TO HILBERT SPACE VALUED SDES

In this section, we establish existence and uniqueness of (local) strong solutions to Hilbert space valued SDEs of the type (1.4).

Let \mathcal{H} be a separable Hilbert space and let $B \in \mathcal{E}$ be a set with $F(B^c) < \infty$. Furthermore, let $a : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow \mathcal{H}$ and $b : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow L_2^0(\mathcal{H})$ be $\mathcal{P} \otimes \mathcal{B}(\mathcal{H})$ -measurable mappings, and let $c : \Omega \times \mathbb{R}_+ \times \mathcal{H} \times E \rightarrow \mathcal{H}$ be a $\mathcal{P} \otimes \mathcal{B}(\mathcal{H}) \otimes \mathcal{E}$ -measurable mapping.

3.1. Definition. *We say that existence of (local) strong solutions to (1.4) holds, if for each \mathcal{F}_0 -measurable random variable $y_0 : \Omega \rightarrow \mathcal{H}$ there exists a (local) strong solution to (1.4) with initial condition y_0 (and some strictly positive lifetime $\tau > 0$).*

3.2. Definition. *We say that uniqueness of (local) strong solutions to (1.4) holds, if for two (local) strong solutions to (1.4) with initial conditions y_0 and y'_0 (and lifetimes τ and τ') we have up to indistinguishability*

$$\begin{aligned} Y \mathbb{1}_{\{y_0=y'_0\}} &= Y' \mathbb{1}_{\{y_0=y'_0\}} \\ (Y^{\tau \wedge \tau'} \mathbb{1}_{\{y_0=y'_0\}} &= (Y')^{\tau \wedge \tau'} \mathbb{1}_{\{y_0=y'_0\}}). \end{aligned}$$

Note that uniqueness of local strong solutions to (1.4) implies uniqueness of strong solutions to (1.4). This is seen by setting $\tau := \infty$ and $\tau' := \infty$.

3.3. Definition. *We say that the mappings $(a, b, c|_B)$ are locally Lipschitz, if \mathbb{P} -almost surely*

$$\left(\int_B \|c(t, y, x)\|^2 F(dx) \right)^{1/2} < \infty \quad \text{for all } t \in \mathbb{R}_+ \text{ and all } y \in \mathcal{H},$$

and for each $n \in \mathbb{N}$ there is a non-decreasing function $L_n : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that \mathbb{P} -almost surely

$$(3.1) \quad \|a(t, y_1) - a(t, y_2)\| \leq L_n(t) \|y_1 - y_2\|,$$

$$(3.2) \quad \|b(t, y_1) - b(t, y_2)\|_{L_2^0(\mathcal{H})} \leq L_n(t) \|y_1 - y_2\|,$$

$$(3.3) \quad \left(\int_B \|c(t, y_1, x) - c(t, y_2, x)\|^2 F(dx) \right)^{1/2} \leq L_n(t) \|y_1 - y_2\|$$

for all $t \in \mathbb{R}_+$ and all $y_1, y_2 \in \mathcal{H}$ with $\|y_1\|, \|y_2\| \leq n$.

3.4. Definition. *We say that the mappings $(a, b, c|_B)$ satisfy the linear growth condition, if there exists a non-decreasing function $K : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that \mathbb{P} -almost surely*

$$(3.4) \quad \|a(t, y)\| \leq K(t)(1 + \|y\|),$$

$$(3.5) \quad \|b(t, y)\|_{L_2^0(\mathcal{H})} \leq K(t)(1 + \|y\|),$$

$$(3.6) \quad \left(\int_B \|c(t, y, x)\|^2 F(dx) \right)^{1/2} \leq K(t)(1 + \|y\|)$$

for all $t \in \mathbb{R}_+$ and all $y \in \mathcal{H}$.

3.5. Definition. *We say that the mappings $(a, b, c|_B)$ are locally bounded, if for each $n \in \mathbb{N}$ there is a non-decreasing function $M_n : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that \mathbb{P} -almost*

surely

$$\begin{aligned} \|a(t, y)\| &\leq M_n(t), \\ \|b(t, y)\|_{L^0_2(H)} &\leq M_n(t), \\ \left(\int_B \|c(t, y, x)\|^2 F(dx) \right)^{1/2} &\leq M_n(t) \end{aligned}$$

for all $t \in \mathbb{R}_+$ and all $y \in \mathcal{H}$ with $\|y\| \leq n$.

For a finite stopping time τ and a set $\Gamma \in \mathcal{F}_\tau$ we define the mappings $a^{(\tau, \Gamma)} : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow \mathcal{H}$, $b^{(\tau, \Gamma)} : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow L^0_2(\mathcal{H})$ and $c^{(\tau, \Gamma)} : \Omega \times \mathbb{R}_+ \times \mathcal{H} \times E \rightarrow \mathcal{H}$ as

$$(3.7) \quad a^{(\tau, \Gamma)}(t, y) := a(\tau + t, y) \mathbb{1}_\Gamma,$$

$$(3.8) \quad b^{(\tau, \Gamma)}(t, y) := b(\tau + t, y) \mathbb{1}_\Gamma,$$

$$(3.9) \quad c^{(\tau, \Gamma)}(t, y, x) := c(\tau + t, y, x) \mathbb{1}_\Gamma.$$

By Lemma 2.2, the mappings $a^{(\tau, \Gamma)}$ and $b^{(\tau, \Gamma)}$ are $\mathcal{P}^{(\tau)} \otimes \mathcal{B}(\mathcal{H})$ -measurable, and $c^{(\tau, \Gamma)}$ is $\mathcal{P}^{(\tau)} \otimes \mathcal{B}(\mathcal{H}) \otimes \mathcal{E}$ -measurable. We shall also use the notation

$$(3.10) \quad a^{(\tau)} := a^{(\tau, \Omega)}, \quad b^{(\tau)} := b^{(\tau, \Omega)} \quad \text{and} \quad c^{(\tau)} := c^{(\tau, \Omega)}.$$

3.6. Lemma. *Suppose that $\tau \mathbb{1}_\Gamma$ is bounded. Then, the following statements are true:*

- (1) *If $(a, b, c|_B)$ are locally Lipschitz, then $(a^{(\tau, \Gamma)}, b^{(\tau, \Gamma)}, c^{(\tau, \Gamma)}|_B)$ are locally Lipschitz, too.*
- (2) *If $(a, b, c|_B)$ satisfy the linear growth condition, then $(a^{(\tau, \Gamma)}, b^{(\tau, \Gamma)}, c^{(\tau, \Gamma)}|_B)$ satisfy the linear growth condition, too.*

Proof. Suppose that $(a, b, c|_B)$ satisfy the linear growth condition. Since $\tau \mathbb{1}_\Gamma$ is bounded, there exists a constant $T \geq 0$ such that $\tau \mathbb{1}_\Gamma \leq T$. The mapping $\tilde{K} := K(\bullet + T) : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is non-decreasing, and we have \mathbb{P} -almost surely

$$\|a^{(\tau, \Gamma)}(t, y)\| = \|a(t + \tau, y) \mathbb{1}_\Gamma\| \leq K(t + \tau) \mathbb{1}_\Gamma (1 + \|y\|) \leq \tilde{K}(t) (1 + \|y\|)$$

for all $t \in \mathbb{R}_+$ and $y \in \mathcal{H}$. Analogous estimates for $b^{(\tau, \Gamma)}$ and $c^{(\tau, \Gamma)}$ prove that $(a^{(\tau, \Gamma)}, b^{(\tau, \Gamma)}, c^{(\tau, \Gamma)}|_B)$ satisfy the linear growth condition, too. The remaining statement is proven analogously. \square

3.7. Lemma. *Let τ and ϱ be two finite stopping times and let $\Gamma \in \mathcal{F}_\tau$ be a set with $\Gamma \subset \{\tau \leq \varrho\}$. If Y is a \mathbb{F} -adapted local strong solution to (1.4) with lifetime ϱ , then*

$$(3.11) \quad Y^{(\tau, \Gamma)} := Y_{\tau+\bullet} \mathbb{1}_\Gamma$$

is a $\mathbb{F}^{(\tau)}$ -adapted local strong solution to (1.4) with parameters

$$(3.12) \quad a = a^{(\tau, \Gamma)}, \quad b = b^{(\tau, \Gamma)}, \quad c = c^{(\tau, \Gamma)}, \quad W = W^{(\tau)}, \quad \mu = \mu^{(\tau)},$$

initial condition $Y_\tau \mathbb{1}_\Gamma$, and lifetime $(\varrho - \tau)^+$.

Proof. The process $Y^{(\tau, \Gamma)}$ given by (3.11) is $\mathbb{F}^{(\tau)}$ -adapted, and we have

$$\begin{aligned} Y_t^{(\tau, \Gamma)} \mathbb{1}_{[0, (\varrho - \tau)^+]}(t) &= Y_{\tau+t} \mathbb{1}_\Gamma \mathbb{1}_{[0, (\varrho - \tau)^+]}(t) = [Y_\tau + (Y_{\tau+t} - Y_\tau)] \mathbb{1}_\Gamma \mathbb{1}_{[0, (\varrho - \tau)^+]}(t) \\ &= \left[Y_\tau + \int_\tau^{\tau+t} a(s, Y_s) ds + \int_\tau^{\tau+t} b(s, Y_s) dW_s \right. \\ &\quad + \int_\tau^{\tau+t} \int_B c(s, Y_{s-}, x) (\mu(ds, dx) - F(dx) ds) \\ &\quad \left. + \int_\tau^{\tau+t} \int_{B^c} c(s, Y_{s-}, x) \mu(ds, dx) \right] \mathbb{1}_\Gamma \mathbb{1}_{[0, (\varrho - \tau)^+]}(t). \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} Y_t^{(\tau, \Gamma)} \mathbb{1}_{[0, (\varrho - \tau)^+]}(t) &= \left[Y_\tau \mathbb{1}_\Gamma + \int_0^t a(\tau + s, Y_{\tau+s}) \mathbb{1}_\Gamma ds + \int_0^t b(\tau + s, Y_{\tau+s}) \mathbb{1}_\Gamma dW_s^{(\tau)} \right. \\ &\quad + \int_0^t \int_B c(\tau + s, Y_{(\tau+s)^-}, x) \mathbb{1}_\Gamma (\mu^{(\tau)}(ds, dx) - F(dx)ds) \\ &\quad \left. + \int_0^t \int_{B^c} c(\tau + s, Y_{(\tau+s)^-}, x) \mathbb{1}_\Gamma \mu^{(\tau)}(ds, dx) \right] \mathbb{1}_{[0, (\varrho - \tau)^+]}(t). \end{aligned}$$

Taking into account the Definitions (3.7)–(3.9) of $a^{(\tau, \Gamma)}$, $b^{(\tau, \Gamma)}$, $c^{(\tau, \Gamma)}$ and the Definition (3.11) of $Y^{(\tau, \Gamma)}$, it follows that

$$\begin{aligned} Y_t^{(\tau, \Gamma)} \mathbb{1}_{[0, (\varrho - \tau)^+]}(t) &= \left[Y_\tau \mathbb{1}_\Gamma + \int_0^t a^{(\tau, \Gamma)}(s, Y_s^{(\tau, \Gamma)}) ds + \int_0^t b^{(\tau, \Gamma)}(s, Y_s^{(\tau, \Gamma)}) dW_s^{(\tau)} \right. \\ &\quad + \int_0^t \int_B c^{(\tau)}(s, Y_{s^-}^{(\tau, \Gamma)}, x) (\mu^{(\tau)}(ds, dx) - F(dx)ds) \\ &\quad \left. + \int_0^t \int_{B^c} c^{(\tau)}(s, Y_{s^-}^{(\tau, \Gamma)}, x) \mu^{(\tau)}(ds, dx) \right] \mathbb{1}_{[0, (\varrho - \tau)^+]}(t). \end{aligned}$$

Consequently, $Y^{(\tau, \Gamma)}$ is a local strong solution to (1.4) with parameters (3.12), initial condition $Y_\tau \mathbb{1}_\Gamma$, and lifetime $(\varrho - \tau)^+$. \square

3.8. Lemma. *Let $\tau \leq \varrho$ be two finite stopping times. If $Y^{(0)}$ is a \mathbb{F} -adapted local strong solution to (1.4) with lifetime τ , and $Y^{(\tau)}$ is a $\mathbb{F}^{(\tau)}$ -adapted local strong solution to (1.4) with parameters*

$$(3.13) \quad a = a^{(\tau)}, \quad b = b^{(\tau)}, \quad c = c^{(\tau)}, \quad W = W^{(\tau)}, \quad \mu = \mu^{(\tau)},$$

initial condition $Y_\tau^{(0)}$, and lifetime $\varrho - \tau$, then

$$(3.14) \quad Y := Y^{(0)} \mathbb{1}_{[0, \tau]} + Y_{\bullet - \tau}^{(\tau)} \mathbb{1}_{[\tau, \varrho]}$$

is a \mathbb{F} -adapted local strong solution to (1.4) with lifetime ϱ .

Proof. Let $t \in \mathbb{R}_+$ be arbitrary. Then, the random variable $Y_t^{(0)} \mathbb{1}_{\{\tau \geq t\}}$ is \mathcal{F}_t -measurable. Let $C \in \mathcal{B}(\mathcal{H})$ be an arbitrary Borel set. We define $D_C \in \mathcal{F}_t$ as

$$D_C := \begin{cases} (\{\tau < t\} \cap \{t \leq \varrho\})^c, & \text{if } 0 \in C, \\ \emptyset, & \text{if } 0 \notin C. \end{cases}$$

According to Lemma 2.1, the mapping $(t - \tau)^+$ is a $\mathbb{F}^{(\tau)}$ -stopping time. Therefore, we get

$$\{Y_{(t-\tau)^+}^{(\tau)} \in C\} \in \mathcal{F}_{(t-\tau)^+}^{(\tau)} = \mathcal{F}_{\tau+(t-\tau)^+},$$

and hence, we obtain

$$\begin{aligned} \{Y_{t-\tau}^{(\tau)} \mathbb{1}_{\{\tau < t \leq \varrho\}} \in C\} &= \{Y_{(t-\tau)^+}^{(\tau)} \mathbb{1}_{\{\tau < t \leq \varrho\}} \in C\} \\ &= (\{\tau < t\} \cap \{t \leq \varrho\} \cap \{Y_{(t-\tau)^+}^{(\tau)} \in C\}) \cup D_C \\ &= (\{\varrho \geq t\} \cap \{\tau \neq t\} \cap \{\tau \leq t\} \cap \{Y_{(t-\tau)^+}^{(\tau)} \in C\}) \cup D_C \\ &= (\{\varrho \geq t\} \cap \{\tau \neq t\} \cap \{\tau + (t - \tau)^+ = t\} \cap \{Y_{(t-\tau)^+}^{(\tau)} \in C\}) \cup D_C \in \mathcal{F}_t, \end{aligned}$$

showing that the process Y defined in (3.14) is \mathbb{F} -adapted. Moreover, since $Y^{(\tau)}$ is local strong solution to (1.4) with initial condition $Y_\tau^{(0)}$ and lifetime $\varrho - \tau$, we have

$$\begin{aligned} Y_{t-\tau}^{(\tau)} \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t) &= \left[Y_\tau^{(0)} + \int_0^{t-\tau} a^{(\tau)}(s, Y_s^{(\tau)}) ds + \int_0^{t-\tau} b^{(\tau)}(s, Y_s^{(\tau)}) dW_s^{(\tau)} \right. \\ &\quad + \int_0^{t-\tau} \int_B c^{(\tau)}(s, Y_{s-}^{(\tau)}, x) (\mu^{(\tau)}(ds, dx) - F(dx) ds) \\ &\quad \left. + \int_0^{t-\tau} \int_{B^c} c^{(\tau)}(s, Y_{s-}^{(\tau)}, x) \mu^{(\tau)}(ds, dx) \right] \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t). \end{aligned}$$

By the Definitions (3.7)–(3.10) of $a^{(\tau)}$, $b^{(\tau)}$, $c^{(\tau)}$, we obtain

$$\begin{aligned} Y_{t-\tau}^{(\tau)} \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t) &= \left[Y_\tau^{(0)} + \int_0^{t-\tau} a(\tau + s, Y_s^{(\tau)}) ds + \int_0^{t-\tau} b(\tau + s, Y_s^{(\tau)}) dW_s^{(\tau)} \right. \\ &\quad + \int_0^{t-\tau} \int_B c(\tau + s, Y_{s-}^{(\tau)}, x) (\mu^{(\tau)}(ds, dx) - F(dx) ds) \\ &\quad \left. + \int_0^{t-\tau} \int_{B^c} c(\tau + s, Y_{s-}^{(\tau)}, x) \mu^{(\tau)}(ds, dx) \right] \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t). \end{aligned}$$

Therefore, we get

$$\begin{aligned} Y_{t-\tau}^{(\tau)} \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t) &= \left[Y_\tau^{(0)} + \int_\tau^t a(s, Y_{s-\tau}^{(\tau)}) ds + \int_\tau^t b(s, Y_{s-\tau}^{(\tau)}) dW_s \right. \\ &\quad + \int_\tau^t \int_B c(s, Y_{(s-\tau)-}^{(\tau)}, x) (\mu(ds, dx) - F(dx) ds) \\ &\quad \left. + \int_\tau^t \int_{B^c} c(s, Y_{(s-\tau)-}^{(\tau)}, x) \mu(ds, dx) \right] \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t). \end{aligned}$$

By the Definition (3.14) of Y we obtain

$$\begin{aligned} Y_{t-\tau}^{(\tau)} \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t) &= \left[Y_\tau^{(0)} + \int_\tau^t a(s, Y_s) ds + \int_\tau^t b(s, Y_s) dW_s \right. \\ &\quad + \int_\tau^t \int_B c(s, Y_{s-}, x) (\mu(ds, dx) - F(dx) ds) \\ &\quad \left. + \int_\tau^t \int_{B^c} c(s, Y_{s-}, x) \mu(ds, dx) \right] \mathbb{1}_{\llbracket \tau, \varrho \rrbracket}(t). \end{aligned}$$

Since $Y^{(0)}$ is a local strong solution to (1.4) with lifetime τ , we deduce that the process Y given by (3.14) is a local strong solution to (1.4) with lifetime ϱ . \square

Let $k \in \mathbb{N}_0$ be arbitrary. By Lemmas 2.1 and 2.3, the mapping $\varrho_{k+1} - \varrho_k$ is a strictly positive $\mathbb{F}^{(\varrho_k)}$ -stopping time. Furthermore, let $\Gamma \in \mathcal{F}_{\varrho_k}$ be arbitrary and let $y_0^{(\varrho_k)} : \Omega \rightarrow \mathcal{H}$ be an arbitrary $\mathcal{F}_0^{(\varrho_k)}$ -measurable random variable.

3.9. Lemma. *If $Y^{(\varrho_k, \Gamma)}$ is a $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solution to (1.4) with parameters*

$$(3.15) \quad a = a^{(\varrho_k, \Gamma)}, \quad b = b^{(\varrho_k, \Gamma)}, \quad c = c^{(\varrho_k, \Gamma)}, \quad W = W^{(\varrho_k)}, \quad \mu = \mu^{(\varrho_k)},$$

initial condition $y_0^{(\varrho_k)} \mathbb{1}_\Gamma$, and lifetime τ , then

$$(3.16) \quad Y^{(\varrho_k, \Gamma)-} := Y^{(\varrho_k, \Gamma)} - c(\varrho_{k+1}, Y_{(\varrho_{k+1}-\varrho_k)-}^{(\varrho_k, \Gamma)}, \xi_{\varrho_{k+1}}) \mathbb{1}_{\llbracket \varrho_{k+1}-\varrho_k \rrbracket} \mathbb{1}_{\{\varrho_{k+1}-\varrho_k \leq \tau\}} \mathbb{1}_\Gamma$$

is a $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solution to (1.5) with parameters (3.15), initial condition $y_0^{(\varrho_k)} \mathbb{1}_\Gamma$, and lifetime $\tau \wedge (\varrho_{k+1} - \varrho_k)$.

Proof. We define $J : \Omega \rightarrow \mathcal{H}$ as

$$J := c(\varrho_{k+1}, Y_{(\varrho_{k+1}-\varrho_k)^-}^{(\varrho_k, \Gamma)}, \xi_{\varrho_{k+1}}) \mathbb{1}_{\{\varrho_{k+1}-\varrho_k \leq \tau\}} \mathbb{1}_\Gamma$$

and the stochastic process $(J_t)_{t \geq 0}$ as $J_t := J \mathbb{1}_{[\varrho_{k+1}-\varrho_k]}(t)$. By Lemma 2.2, the mapping J is $\mathcal{F}_{\varrho_{k+1}}$ -measurable. Let $C \in \mathcal{B}(\mathcal{H})$ be an arbitrary Borel set. We define $D_C \in \mathcal{F}_t$ as

$$D_C := \begin{cases} \{\varrho_{k+1} - \varrho_k \neq t\}, & \text{if } 0 \in C, \\ \emptyset, & \text{if } 0 \notin C. \end{cases}$$

Then, for each $t \in \mathbb{R}_+$ we have

$$\begin{aligned} \{J_t \in C\} &= \{J \mathbb{1}_{[\varrho_{k+1}-\varrho_k]}(t) \in C\} = (\{J \in C\} \cap \{\varrho_{k+1} - \varrho_k = t\}) \cup D_C \\ &= (\{J \in C\} \cap \{\varrho_{k+1} = \varrho_k + t\}) \cup D_C \in \mathcal{F}_{\varrho_k+t} = \mathcal{F}_t^{(\varrho_k)}. \end{aligned}$$

Consequently, the process $Y^{(\varrho_k, \Gamma)^-}$ defined in (3.16) is $\mathbb{F}^{(\varrho_k)}$ -adapted. Furthermore, by the Definition (3.16) we have

$$Y_-^{(\varrho_k, \Gamma)} \mathbb{1}_{[0, \tau \wedge (\varrho_{k+1}-\varrho_k)]} = Y_-^{(\varrho_k, \Gamma)^-} \mathbb{1}_{[0, \tau \wedge (\varrho_{k+1}-\varrho_k)]}$$

and, by the Definition (3.9) of $c^{(\varrho_k, \Gamma)}$ and identity (2.1) we obtain

$$\begin{aligned} & \left(\int_0^t \int_{B^c} c^{(\varrho_k, \Gamma)}(s, Y_{s^-}^{(\varrho_k, \Gamma)}, x) \mu^{(\varrho_k)}(ds, dx) \right) \mathbb{1}_{[0, \tau \wedge (\varrho_{k+1}-\varrho_k)]}(t) \\ &= \left(\int_0^t \int_{B^c} c(\varrho_k + s, Y_{s^-}^{(\varrho_k, \Gamma)}, x) \mathbb{1}_\Gamma \mu^{(\varrho_k)}(ds, dx) \right) \mathbb{1}_{[0, \tau \wedge (\varrho_{k+1}-\varrho_k)]}(t) \\ &= \left(\int_{\varrho_k}^{\varrho_k+t} \int_{B^c} c(s, Y_{(s-\varrho_k)^-}^{(\varrho_k, \Gamma)}, x) \mathbb{1}_\Gamma \mu(ds, dx) \right) \mathbb{1}_{[0, \tau \wedge (\varrho_{k+1}-\varrho_k)]}(t) \\ &= \left(\sum_{n \in \mathbb{N}} c(\kappa_n, Y_{(\kappa_n-\varrho_k)^-}^{(\varrho_k, \Gamma)}, \xi_{\kappa_n}) \mathbb{1}_{\{\xi_{\kappa_n} \notin B\}} \mathbb{1}_{\{\varrho_k < \kappa_n \leq \varrho_k+t\}} \right) \mathbb{1}_{[0, \tau \wedge (\varrho_{k+1}-\varrho_k)]}(t) \mathbb{1}_\Gamma \\ &= c(\varrho_{k+1}, Y_{(\varrho_{k+1}-\varrho_k)^-}^{(\varrho_k, \Gamma)}, \xi_{\varrho_{k+1}}) \mathbb{1}_{[\varrho_{k+1}-\varrho_k]}(t) \mathbb{1}_{\{\varrho_{k+1}-\varrho_k \leq \tau\}} \mathbb{1}_\Gamma, \end{aligned}$$

showing that $Y^{(\varrho_k, \Gamma)^-}$ is a local strong solution to (1.5) with parameters (3.15) and lifetime $\tau \wedge (\varrho_{k+1} - \varrho_k)$. \square

3.10. Lemma. *If $Y^{(\varrho_k, \Gamma)^-}$ is a $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solution to (1.5) with parameters (3.15), initial condition $y_0^{(\varrho_k)} \mathbb{1}_\Gamma$, and lifetime τ , then*

$$(3.17) \quad Y^{(\varrho_k, \Gamma)} := Y^{(\varrho_k, \Gamma)^-} + c(\varrho_{k+1}, Y_{(\varrho_{k+1}-\varrho_k)^-}^{(\varrho_k, \Gamma)^-}, \xi_{\varrho_{k+1}}) \mathbb{1}_{[\varrho_{k+1}-\varrho_k]} \mathbb{1}_{\{\varrho_{k+1}-\varrho_k \leq \tau\}} \mathbb{1}_\Gamma$$

is a $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solution to (1.4) with parameters (3.15), initial condition $y_0^{(\varrho_k)} \mathbb{1}_\Gamma$, and lifetime $\tau \wedge (\varrho_{k+1} - \varrho_k)$.

Proof. The proof is analogous to that of Lemma 3.9. \square

3.1. Uniqueness of strong solutions to Hilbert space valued SDEs. Now, we shall deal with the uniqueness of strong solutions to the SDE (1.4).

3.11. Proposition. *We suppose that the mappings $(a, b, c|_B)$ are locally Lipschitz. Then, uniqueness of local strong solutions to (1.5) holds.*

Proof. We can adopt a standard technique (see, e.g. the proof of Theorem 5.2.5 in [15]), where we apply the Itô isometry and Gronwall's lemma. \square

3.12. Theorem. *We suppose that the mappings $(a, b, c|_B)$ are locally Lipschitz. Then, uniqueness of local strong solutions to (1.4) holds.*

Proof. Let Y and Y' be two local strong solutions to (1.5) with initial conditions y_0 and y'_0 , and lifetimes τ and τ' . By induction, we will prove that up to indistinguishability

$$(3.18) \quad Y \mathbb{1}_{[0, \tau \wedge \tau' \wedge \varrho_k]} \mathbb{1}_{\{y_0 = y'_0\}} = Y' \mathbb{1}_{[0, \tau \wedge \tau' \wedge \varrho_k]} \mathbb{1}_{\{y_0 = y'_0\}} \quad \text{for all } k \in \mathbb{N}_0.$$

The identity (3.18) holds true for $k = 0$, because by Lemma 2.3 we have $\varrho_0 = 0$.

For the induction step $k \rightarrow k + 1$ we suppose that identity (3.18) is satisfied. We define the stopping time $\tau_k := \tau \wedge \tau' \wedge \varrho_{k+1}$ and the set $\Gamma_k := \{\varrho_k \leq \tau_k\} \cap \{y_0 = y'_0\} \in \mathcal{F}_{\varrho_k}$. By Lemma 3.7, the processes $Y^{(\varrho_k, \Gamma_k)} := Y_{\varrho_k + \bullet} \mathbb{1}_{\Gamma_k}$ and $Y'^{(\varrho_k, \Gamma_k)} := Y'_{\varrho_k + \bullet} \mathbb{1}_{\Gamma_k}$ defined according to (3.11) are $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solutions to (1.4) with parameters (3.12), where $\tau = \varrho_k$ and $\Gamma = \Gamma_k$, initial conditions $Y_{\varrho_k} \mathbb{1}_{\Gamma_k}$ and $Y'_{\varrho_k} \mathbb{1}_{\Gamma_k}$, and lifetime $(\tau_k - \varrho_k)^+$.

Let $n \in \mathbb{N}$ be arbitrary and set $\Gamma_{kn} := \Gamma_k \cap \{\varrho_k \leq n\} \in \mathcal{F}_{\varrho_k}$. The processes $Y^{(\varrho_k, \Gamma_{kn})} := Y^{(\varrho_k, \Gamma_k)} \mathbb{1}_{\Gamma_{kn}}$ and $Y'^{(\varrho_k, \Gamma_{kn})} := Y'^{(\varrho_k, \Gamma_k)} \mathbb{1}_{\Gamma_{kn}}$ are $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solutions to (1.4) with parameters (3.15), where $\Gamma = \Gamma_{kn}$, initial conditions $Y_{\varrho_k} \mathbb{1}_{\Gamma_{kn}}$ and $Y'_{\varrho_k} \mathbb{1}_{\Gamma_{kn}}$, and lifetime $(\tau_k - \varrho_k)^+$. By Lemma 3.9, the processes $Y^{(\varrho_k, \Gamma_{kn})-}$ and $Y'^{(\varrho_k, \Gamma_{kn})-}$ defined according to (3.16) are $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solutions to (1.5) with parameters (3.15), where $\Gamma = \Gamma_{kn}$, initial conditions $Y_{\varrho_k} \mathbb{1}_{\Gamma_{kn}}$ and $Y'_{\varrho_k} \mathbb{1}_{\Gamma_{kn}}$, and lifetime $(\tau_k - \varrho_k)^+$. According to Lemma 3.6, the mappings $(a^{(\varrho_k, \Gamma_{kn})}, b^{(\varrho_k, \Gamma_{kn})}, c^{(\varrho_k, \Gamma_{kn})})|_B$ are locally Lipschitz, too. Therefore, by Proposition 3.11 we have up to indistinguishability

$$Y^{(\varrho_k, \Gamma_{kn})-} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} = Y'^{(\varrho_k, \Gamma_{kn})-} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} \quad \text{for all } n \in \mathbb{N}.$$

By the Definition (3.16), we deduce that up to indistinguishability

$$Y^{(\varrho_k, \Gamma_{kn})} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} = Y'^{(\varrho_k, \Gamma_{kn})} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} \quad \text{for all } n \in \mathbb{N},$$

and hence, we have up to indistinguishability

$$Y^{(\varrho_k, \Gamma_k)} \mathbb{1}_{\{\varrho_k \leq n\}} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} = Y'^{(\varrho_k, \Gamma_k)} \mathbb{1}_{\{\varrho_k \leq n\}} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} \quad \text{for all } n \in \mathbb{N}.$$

By Lemma 2.3 we have $\mathbb{P}(\varrho_k < \infty) = 1$, and hence, we get up to indistinguishability

$$Y^{(\varrho_k, \Gamma_k)} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} = Y'^{(\varrho_k, \Gamma_k)} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]}.$$

Therefore, we have up to indistinguishability

$$Y_{\varrho_k + \bullet} \mathbb{1}_{\{\varrho_k \leq \tau_k\}} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} \mathbb{1}_{\{y_0 = y'_0\}} = Y'_{\varrho_k + \bullet} \mathbb{1}_{\{\varrho_k \leq \tau_k\}} \mathbb{1}_{[0, (\tau_k - \varrho_k)^+]} \mathbb{1}_{\{y_0 = y'_0\}}.$$

Consequently, we have up to indistinguishability

$$Y \mathbb{1}_{\{\varrho_k \leq \tau_k\}} \mathbb{1}_{[\varrho_k, \tau_k]} \mathbb{1}_{\{y_0 = y'_0\}} = Y' \mathbb{1}_{\{\varrho_k \leq \tau_k\}} \mathbb{1}_{[\varrho_k, \tau_k]} \mathbb{1}_{\{y_0 = y'_0\}}.$$

Together with the induction hypothesis, it follows that

$$Y \mathbb{1}_{[0, \tau_k]} \mathbb{1}_{\{y_0 = y'_0\}} = Y' \mathbb{1}_{[0, \tau_k]} \mathbb{1}_{\{y_0 = y'_0\}},$$

which establishes (3.18). Since by Lemma 2.3 we have $\mathbb{P}(\varrho_k \rightarrow \infty) = 1$, we deduce

$$Y^{\tau \wedge \tau'} \mathbb{1}_{\{y_0 = y'_0\}} = (Y')^{\tau \wedge \tau'} \mathbb{1}_{\{y_0 = y'_0\}},$$

completing the proof. \square

3.2. Existence of strong solutions to Hilbert space valued SDEs. Now, we shall deal with the existence of strong solutions to the SDE (1.4).

3.13. Proposition. *We suppose that the mappings $(a, b, c|_B)$ are locally Lipschitz and satisfy the linear growth condition. Then, existence of strong solutions to (1.5) holds.*

Proof. If the mappings $(a, b, c|_B)$ are Lipschitz continuous, then we have existence and uniqueness of strong solutions to (1.5) for every initial condition $y_0 \in \mathcal{L}^2(\mathcal{F}_0; \mathcal{H})$, see, e.g. [7, Cor. 10.3].

For $(a, b, c|_B)$ being locally Lipschitz and satisfying the linear growth condition, for any initial condition $y_0 \in \mathcal{L}^2(\mathcal{F}_0; \mathcal{H})$ we adopt the technique from the proof of [19, Thm. 4.11]. For $k \in \mathbb{N}$ we define the retraction

$$R_k : \mathcal{H} \rightarrow \mathcal{H}, \quad R_k(y) := \begin{cases} y, & \text{if } \|y\| \leq k, \\ k \frac{y}{\|y\|}, & \text{if } \|y\| > k, \end{cases}$$

and the mappings $a_k : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow \mathcal{H}$, $b_k : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow L^0_2(\mathcal{H})$ and $c_k : \Omega \times \mathbb{R}_+ \times \mathcal{H} \times E \rightarrow \mathcal{H}$ as

$$a_k := a \circ R_k, \quad b_k := b \circ R_k \quad \text{and} \quad c_k(\bullet, x) := c(\bullet, x) \circ R_k.$$

These mappings are Lipschitz continuous, and hence there exists a strong solution $Y^{(k)}$ to the SDE (1.5) with parameters $a = a_k$, $b = b_k$ and $c = c_k$, and initial condition y_0 . Using the linear growth condition, Gronwall's lemma and Doob's martingale inequality, we can show that $\mathbb{P}(\tau_k \rightarrow \infty) = 1$, where

$$\tau_k := \inf\{t \geq 0 : \|Y_t^{(k)}\| > k\}, \quad k \in \mathbb{N}_0,$$

i.e. the solutions do not explode. Consequently, the process

$$Y := y_0 \mathbb{1}_{[\tau_0]} + \sum_{k \in \mathbb{N}} Y^{(k)} \mathbb{1}_{[\tau_{k-1}, \tau_k]}$$

is a strong solution to (1.5) with initial condition y_0 .

Finally, for a general \mathcal{F}_0 -measurable initial condition $y_0 : \Omega \rightarrow \mathcal{H}$, the process $Y := \sum_{k \in \mathbb{N}} Y^{(k)} \mathbb{1}_{\Omega_k}$ is a strong solution to (1.5) with initial condition y_0 , where $(\Omega_k)_{k \in \mathbb{N}} \subset \mathcal{F}_0$ denotes the partition of Ω given by $\Omega_k := \{\|y_0\| \in [k-1, k)\}$, and where for each $k \in \mathbb{N}$ the process $Y^{(k)}$ denotes a strong solution to (1.5) with initial condition $y_0 \mathbb{1}_{\Omega_k}$. \square

3.14. Theorem. *We suppose that the mappings $(a, b, c|_B)$ are locally Lipschitz and satisfy the linear growth condition. Then, existence of strong solutions to (1.4) holds.*

Proof. Let $y_0 : \Omega \rightarrow \mathcal{H}$ be an arbitrary \mathcal{F}_0 -measurable random variable. By induction, we will prove that for each $k \in \mathbb{N}_0$ there exists a local strong solution $Y^{(k)}$ to (1.4) with initial condition y_0 and lifetime ϱ_k . By Lemma 2.3 we have $\varrho_0 = 0$, providing the assertion for $k = 0$.

For the induction step $k \rightarrow k+1$ let $Y^{(k)}$ be a local strong solution to (1.4) with initial condition y_0 and lifetime ϱ_k . Let $n \in \mathbb{N}$ be arbitrary and set $\Gamma_{kn} := \{\varrho_k \in [n-1, n)\} \in \mathcal{F}_{\varrho_k}$. By Lemma 3.6, the mappings $(a^{(\varrho_k, \Gamma_{kn})}, b^{(\varrho_k, \Gamma_{kn})}, c^{(\varrho_k, \Gamma_{kn})}|_B)$ are locally Lipschitz, too. Therefore, by Proposition 3.13 there exists a $\mathbb{F}^{(\varrho_k)}$ -adapted strong solution $Y^{(\varrho_k, \Gamma_{kn})}$ to (1.5) with parameters (3.15), where $\Gamma = \Gamma_{kn}$, and initial condition $Y_{\varrho_k}^{(k)} \mathbb{1}_{\Gamma_{kn}}$. By Lemma 3.10, the process $Y^{(\varrho_k, \Gamma_{kn})}$ defined according to (3.17) is a $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solution to (1.4) with parameters (3.15), where $\Gamma = \Gamma_{kn}$, initial condition $Y_{\varrho_k}^{(k)} \mathbb{1}_{\Gamma_{kn}}$, and lifetime $\varrho_{k+1} - \varrho_k$. Noting that $(\Gamma_{kn})_{n \in \mathbb{N}}$ is a partition of Ω , it follows that $Y^{(\varrho_k)} := \sum_{n \in \mathbb{N}} Y^{(\varrho_k, \Gamma_{kn})}$ is a $\mathbb{F}^{(\varrho_k)}$ -adapted local strong solution to (1.4) with initial condition $Y_{\varrho_k}^{(k)}$ and lifetime $\varrho_{k+1} - \varrho_k$. By Lemma 3.8, the process

$$Y^{(k+1)} := Y^{(k)} \mathbb{1}_{[0, \varrho_k]} + Y_{\bullet - \varrho_k}^{(\varrho_k)} \mathbb{1}_{[\varrho_k, \varrho_{k+1}]}$$

defined according to (3.14) is a \mathbb{F} -adapted local strong solution to (1.4) with initial condition y_0 and lifetime ϱ_{k+1} .

Consequently, for each $k \in \mathbb{N}_0$ there exists a local strong solution $Y^{(k)}$ to (1.4) with initial condition y_0 and lifetime ϱ_k . By Lemma 2.3 we have $\mathbb{P}(\varrho_k \rightarrow \infty) = 1$. Hence, it follows that

$$Y := y_0 \mathbb{1}_{[\varrho_0]} + \sum_{k \in \mathbb{N}} Y^{(k)} \mathbb{1}_{[\varrho_{k-1}, \varrho_k]}$$

is a \mathbb{F} -adapted strong solution to (1.4) with initial condition y_0 . \square

3.15. Theorem. *We suppose that the mappings $(a, b, c|_B)$ are locally Lipschitz and locally bounded. Then, existence of local strong solutions to (1.4) holds.*

Proof. Let $y_0 : \Omega \rightarrow \mathcal{H}$ be an arbitrary \mathcal{F}_0 -measurable random variable. We define the partition $(\Omega_k)_{k \in \mathbb{N}} \subset \mathcal{F}_0$ of Ω by $\Omega_k := \{\|y_0\| \in [k-1, k)\}$. Furthermore, for each $k \in \mathbb{N}$ we define the mappings $a_k : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow \mathcal{H}$, $b_k : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow L_2^0(\mathcal{H})$ and $c_k : \Omega \times \mathbb{R}_+ \times \mathcal{H} \times E \rightarrow \mathcal{H}$ as in the proof of Proposition 3.13. These mappings are locally Lipschitz and satisfy the linear growth condition. By Theorem 3.14, there exists a strong solution $Y^{(k)}$ to (1.4) with parameters $a = a_k$, $b = b_k$ and $c = c_k$, and initial condition $y_0 \mathbb{1}_{\Omega_k}$. The stopping time

$$\tau_k := \inf\{t \geq 0 : \|Y_t^{(k)}\| > k\}$$

is strictly positive, and $Y^{(k)}$ is a local strong solution to (1.4) with initial condition $y_0 \mathbb{1}_{\Omega_k}$ and lifetime τ_k . The stopping time $\tau := \sum_{k \in \mathbb{N}} \tau_k \mathbb{1}_{\Omega_k}$ is strictly positive, and the process $Y := \sum_{k \in \mathbb{N}} Y^{(k)} \mathbb{1}_{\Omega_k}$ is a local strong solution to (1.4) with initial condition y_0 and lifetime τ . \square

3.3. Comparison with the method of successive approximations. So far, our investigations provide the following result concerning existence and uniqueness of global strong solutions to the SDE (1.4).

3.16. Theorem. *If $(a, b, c|_B)$ are locally Lipschitz and satisfy the linear growth condition, then existence and uniqueness of strong solutions to (1.4) holds.*

Proof. This is a direct consequence of Theorems 3.12 and 3.14. \square

Now, we shall provide a comparison with reference [3], where the authors also study Hilbert space valued SDEs of the type (1.4). Their result [3, Theorem 2.1] is based on the method of successive approximations (see also [26, 25]) and considerably goes beyond the classical global Lipschitz conditions. For the sake of simplicity, let us recall the required assumptions in the time-homogeneous Markovian framework. In order to apply [3, Theorem 2.1], for some constant $p \geq 2$ we need the estimate

$$(3.19) \quad \begin{aligned} & \|a(y_1) - a(y_2)\|^p + \|b(y_1) - b(y_2)\|_{L_2^0(\mathcal{H})}^p + \int_B \|c(y_1, x) - c(y_2, y)\|^p F(dx) \\ & + \left(\int_B \|c(y_1, x) - c(y_2, x)\|^2 F(dx) \right)^{p/2} \leq \kappa(\|y_1 - y_2\|^p) \quad \text{for all } y_1, y_2 \in \mathcal{H}, \end{aligned}$$

where $\kappa : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ denotes a continuous, nondecreasing function with $\kappa(0) = 0$, and further conditions, which are precisely stated in [3], must be fulfilled. These conditions are satisfied if κ is a continuous, nondecreasing and concave function such that

$$(3.20) \quad \int_0^\epsilon \frac{1}{\kappa(u)} du = \infty \quad \text{for each } \epsilon > 0.$$

In particular, we may choose $\kappa(u) = u$ for $u \in \mathbb{R}_+$, and consequently, both results, Theorem 3.16 and [3, Theorem 2.1], cover the classical situation, where global Lipschitz conditions are imposed.

However, there are situations where [3, Theorem 2.1] can be applied, while Theorem 3.16 does not apply, and vice versa. For the sake of simplicity, in the following two examples we assume that $\mathcal{H} = \mathbb{R}$ and $b \equiv c \equiv 0$.

3.17. Example. We fix an arbitrary constant $0 < \delta < \exp(-1)$ and define the functions $\kappa, \rho : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ by

$$\kappa(u) := \begin{cases} 0, & u = 0, \\ -u \ln u, & 0 < u < \delta, \\ -\delta \ln \delta - (1 + \ln \delta)(u - \delta) & u \geq \delta, \end{cases}$$

as well as

$$\rho(u) := \begin{cases} 0, & u = 0, \\ u \sqrt{-\ln(u^2)}, & 0 < u < \sqrt{\delta}, \\ \sqrt{-\delta \ln \delta - (1 + \ln \delta)(u^2 - \delta)} & u \geq \sqrt{\delta}, \end{cases}$$

cf. [26, Remark 1]. Let $a : \mathbb{R} \rightarrow \mathbb{R}$ be a mapping such that

$$|a(y_1) - a(y_2)| \leq \rho(|y_1 - y_2|) \quad \text{for all } y_1, y_2 \in \mathbb{R}.$$

Then we have the estimate

$$|a(y_1) - a(y_2)|^2 \leq \kappa(|y_1 - y_2|^2) \quad \text{for all } y_1, y_2 \in \mathbb{R},$$

showing that condition (3.19) with $p = 2$ is satisfied. Moreover, κ is a continuous, nondecreasing, concave function and condition (3.20) is satisfied, because for each $0 < \epsilon < \delta$ we have

$$\int_0^\epsilon \frac{1}{\kappa(u)} du = - \int_0^\epsilon \frac{1}{u \ln u} du = - \ln |\ln u| \Big|_{u=0}^{u=\epsilon} = - \ln |\ln \epsilon| + \lim_{u \rightarrow 0} \ln |\ln u| = \infty.$$

Consequently, [3, Theorem 2.1] applies. However, we have

$$\rho'(u) = \sqrt{-\ln(u^2)} - \frac{1}{\sqrt{-\ln(u^2)}} \quad \text{for } u \in (0, \sqrt{\delta}),$$

and thus $\lim_{u \rightarrow 0} \rho'(u) = \infty$. Therefore, the mapping $a : \mathbb{R} \rightarrow \mathbb{R}$ might fail to be locally Lipschitz, and hence, Theorem 3.16 does not apply.

3.18. Example. Let us define the mapping $a : \mathbb{R} \rightarrow \mathbb{R}$ as follows. For $n \in \mathbb{N}_0$ we define a on the interval $[n, n+1]$ by

$$a(y) := \begin{cases} n, & y \in [n, n+1 - \frac{1}{n+1}], \\ n + (n+1)(y - (n+1 - \frac{1}{n+1})), & y \in [n+1 - \frac{1}{n+1}, n+1]. \end{cases}$$

This defines the mapping $a : \mathbb{R}_+ \rightarrow \mathbb{R}$, which we extend to a mapping $a : \mathbb{R} \rightarrow \mathbb{R}$ by symmetry

$$a(y) := a(-y), \quad y \in \mathbb{R}_-.$$

Then, a is locally Lipschitz and satisfies the linear growth condition, and hence, Theorem 3.16 applies. However, there are no constant $p \geq 2$ and no continuous, nondecreasing function $\kappa : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\kappa(0) = 0$ such that

$$(3.21) \quad |a(y_1) - a(y_2)|^p \leq \kappa(|y_1 - y_2|^p) \quad \text{for all } y_1, y_2 \in \mathbb{R}.$$

Suppose, on the contrary, there exists a continuous, nondecreasing function $\kappa : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\kappa(0) = 0$ fulfilling (3.21). Then we have

$$(3.22) \quad |\kappa(u)| \geq 1 \quad \text{for all } u \in (0, 1].$$

Indeed, let $u \in (0, 1]$ be arbitrary. Then, there exists $n \in \mathbb{N}$ with $\frac{1}{n} \leq u$. Moreover, by the definition of the mapping $a : \mathbb{R} \rightarrow \mathbb{R}$ there are $y_1, y_2 \in \mathbb{R}$ such that

$$|y_1 - y_2| \leq \left(\frac{1}{n}\right)^{1/p} \quad \text{and} \quad |a(y_1) - a(y_2)| = 1.$$

Therefore, using the monotonicity of κ and (3.21) we obtain

$$\kappa(u) \geq \kappa\left(\frac{1}{n}\right) \geq \kappa(|y_1 - y_2|^p) \geq |a(y_1) - a(y_2)|^p = 1,$$

showing (3.22). Now, the continuity of κ yields the contradiction $\kappa(0) \geq 1$. Consequently, condition (3.19) is not satisfied, and thus, we cannot use [3, Theorem 2.1] in this case.

4. EXISTENCE AND UNIQUENESS OF MILD SOLUTIONS TO HILBERT SPACE VALUED SPDES

In this section, we establish existence and uniqueness of (local) mild solutions to Hilbert space valued SPDEs of the type (1.3).

Let H be a separable Hilbert space, let $(S_t)_{t \geq 0}$ be a C_0 -semigroup on H with infinitesimal generator $A : \mathcal{D}(A) \subset H \rightarrow H$, and let $B \in \mathcal{E}$ be a set with $F(B^c) < \infty$. Furthermore, let $\alpha : \Omega \times \mathbb{R}_+ \times H \rightarrow H$ and $\sigma : \Omega \times \mathbb{R}_+ \times H \rightarrow L_2^0(H)$ be $\mathcal{P} \otimes \mathcal{B}(H)$ -measurable mappings, and let $\gamma : \Omega \times \mathbb{R}_+ \times H \times E \rightarrow H$ be a $\mathcal{P} \otimes \mathcal{B}(H) \otimes \mathcal{E}$ -measurable mapping.

Throughout this section, we suppose that there exist another separable Hilbert space \mathcal{H} , a C_0 -group $(U_t)_{t \in \mathbb{R}}$ on \mathcal{H} and continuous linear operators $\ell \in L(H, \mathcal{H})$, $\pi \in L(\mathcal{H}, H)$ such that the diagram

$$\begin{array}{ccc} \mathcal{H} & \xrightarrow{U_t} & \mathcal{H} \\ \uparrow \ell & & \downarrow \pi \\ H & \xrightarrow{S_t} & H \end{array}$$

commutes for every $t \in \mathbb{R}_+$, that is

$$(4.1) \quad \pi U_t \ell = S_t \quad \text{for all } t \in \mathbb{R}_+.$$

4.1. Remark. According to [7, Prop. 8.7], this assumption is satisfied if the semigroup $(S_t)_{t \geq 0}$ is pseudo-contractive (one also uses the notion quasi-contractive), that is, there is a constant $\omega \in \mathbb{R}$ such that

$$\|S_t\| \leq e^{\omega t} \quad \text{for all } t \geq 0.$$

This result relies on the Szőkefalvi-Nagy theorem on unitary dilations (see e.g. [24, Thm. I.8.1], or [5, Sec. 7.2]). In the spirit of [24], the group $(U_t)_{t \in \mathbb{R}}$ is called a dilation of the semigroup $(S_t)_{t \geq 0}$.

4.2. Remark. The Szőkefalvi-Nagy theorem was also utilized in [13, 12] in order to establish results concerning stochastic convolution integrals.

Now, we define the mappings $a : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow \mathcal{H}$, $b : \Omega \times \mathbb{R}_+ \times \mathcal{H} \rightarrow L_2^0(\mathcal{H})$ and $c : \Omega \times \mathbb{R}_+ \times \mathcal{H} \times E \rightarrow \mathcal{H}$ by

$$(4.2) \quad a(t, y) := U_{-t} \ell \alpha(t, \pi U_t y),$$

$$(4.3) \quad b(t, y) := U_{-t} \ell \sigma(t, \pi U_t y),$$

$$(4.4) \quad c(t, y, x) := U_{-t} \ell \gamma(t, \pi U_t y, x).$$

Note that a and b are $\mathcal{P} \otimes \mathcal{B}(\mathcal{H})$ -measurable, and that c is $\mathcal{P} \otimes \mathcal{B}(\mathcal{H}) \otimes \mathcal{E}$ -measurable.

4.3. Lemma. The following statements are true:

- (1) If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz, then $(a, b, c|_B)$ are locally Lipschitz, too.
- (2) If $(\alpha, \sigma, \gamma|_B)$ satisfy the linear growth condition, then $(a, b, c|_B)$ satisfy the linear growth condition, too.
- (3) If $(\alpha, \sigma, \gamma|_B)$ are locally bounded, then $(a, b, c|_B)$ are locally bounded, too.

Proof. All three statements are straightforward to check. \square

4.4. Proposition. Let $z_0 : \Omega \rightarrow H$ be a \mathcal{F}_0 -measurable random variable, and let τ be a stopping time. Then, the following statements are true:

- (1) If Y is a local strong solution to (1.4) with initial condition ℓz_0 and lifetime τ , then $Z := \pi UY$ is a local mild solution to (1.3) with initial condition z_0 and lifetime τ .
- (2) If Z is a local mild solution to (1.3) with initial condition z_0 and lifetime τ , then the process Y defined as

$$(4.5) \quad \begin{aligned} Y_t := & \ell z_0 + \int_0^{t \wedge \tau} U_{-s} \ell \alpha(s, Z_s) ds + \int_0^{t \wedge \tau} U_{-s} \ell \sigma(s, Z_s) dW_s \\ & + \int_0^{t \wedge \tau} \int_B U_{-s} \ell \gamma(s, Z_{s-}, x) (\mu(ds, dx) - F(dx) ds) \\ & + \int_0^{t \wedge \tau} \int_{B^c} U_{-s} \ell \gamma(s, Z_{s-}, x) \mu(ds, dx), \quad t \geq 0 \end{aligned}$$

is a local strong solution to (1.4) with initial condition ℓz_0 and lifetime τ , and we have $Z^\tau = \pi UY^\tau$.

Proof. Let Y be a local strong solution to (1.4) with initial condition ℓz_0 and lifetime τ . Then we have

$$\begin{aligned} Z_{t \wedge \tau} = & \pi U_{t \wedge \tau} Y_{t \wedge \tau} = \pi U_{t \wedge \tau} \left(\ell z_0 + \int_0^{t \wedge \tau} a(s, Y_s) ds + \int_0^{t \wedge \tau} b(s, Y_s) dW_s \right. \\ & \left. + \int_0^{t \wedge \tau} \int_B c(s, Y_{s-}, x) (\mu(ds, dx) - F(dx) ds) + \int_0^{t \wedge \tau} \int_{B^c} c(s, Y_{s-}, x) \mu(ds, dx) \right). \end{aligned}$$

By the Definitions (4.2)–(4.4) of a, b, c we obtain

$$\begin{aligned} Z_{t \wedge \tau} = & \pi U_{t \wedge \tau} \left(\ell z_0 + \int_0^{t \wedge \tau} U_{-s} \ell \alpha(s, \pi U_s Y_s) ds + \int_0^{t \wedge \tau} U_{-s} \ell \sigma(s, \pi U_s Y_s) dW_s \right. \\ & + \int_0^{t \wedge \tau} \int_B U_{-s} \ell \gamma(s, \pi U_s Y_{s-}, x) (\mu(ds, dx) - F(dx) ds) \\ & \left. + \int_0^{t \wedge \tau} \int_{B^c} U_{-s} \ell \gamma(s, \pi U_s Y_{s-}, x) \mu(ds, dx) \right). \end{aligned}$$

Therefore, by (4.1), and since $Z = \pi UY$, we arrive at

$$(4.6) \quad \begin{aligned} Z_{t \wedge \tau} = & S_{t \wedge \tau} z_0 + \int_0^{t \wedge \tau} S_{(t \wedge \tau) - s} \alpha(s, Z_s) ds + \int_0^{t \wedge \tau} S_{(t \wedge \tau) - s} \sigma(s, Z_s) dW_s \\ & + \int_0^{t \wedge \tau} \int_B S_{(t \wedge \tau) - s} \gamma(s, Z_{s-}, x) (\mu(ds, dx) - F(dx) ds) \\ & + \int_0^{t \wedge \tau} \int_{B^c} S_{(t \wedge \tau) - s} \gamma(s, Z_{s-}, x) \mu(ds, dx), \end{aligned}$$

showing that Z is a local mild solution to (1.3) with initial condition z_0 and lifetime τ . This establishes the first statement. Now, let Z be a local mild solution to (1.3) with initial condition z_0 and lifetime τ . Then we have (4.6), and therefore, by (4.1)

and the Definition (4.5) of Y we obtain

$$\begin{aligned} Z_{t \wedge \tau} &= \pi U_{t \wedge \tau} \left(\ell z_0 + \int_0^{t \wedge \tau} U_{-s} \ell \alpha(s, Z_s) ds + \int_0^{t \wedge \tau} U_{-s} \ell \sigma(s, Z_s) dW_s \right. \\ &\quad + \int_0^{t \wedge \tau} \int_B U_{-s} \ell \gamma(s, Z_{s-}, x) (\mu(ds, dx) - F(dx) ds) \\ &\quad \left. + \int_0^{t \wedge \tau} \int_{B^c} U_{-s} \ell \gamma(s, Z_{s-}, x) \mu(ds, dx) \right) = \pi U_{t \wedge \tau} Y_{t \wedge \tau}, \end{aligned}$$

showing that $Z^\tau = \pi U Y^\tau$. Therefore, by the Definition (4.5) of Y we obtain

$$\begin{aligned} Y_{t \wedge \tau} &= \ell z_0 + \int_0^{t \wedge \tau} U_{-s} \ell \alpha(s, \pi U_s Y_s) ds + \int_0^{t \wedge \tau} U_{-s} \ell \sigma(s, \pi U_s Y_s) dW_s \\ &\quad + \int_0^{t \wedge \tau} \int_B U_{-s} \ell \gamma(s, \pi U_s Y_{s-}, x) (\mu(ds, dx) - F(dx) ds) \\ &\quad + \int_0^{t \wedge \tau} \int_{B^c} U_{-s} \ell \gamma(s, \pi U_s Y_{s-}, x) \mu(ds, dx). \end{aligned}$$

Taking into account the Definitions (4.2)–(4.4) of a, b, c , we get

$$\begin{aligned} Y_{t \wedge \tau} &= \ell z_0 + \int_0^{t \wedge \tau} a(s, Y_s) ds + \int_0^{t \wedge \tau} b(s, Y_s) dW_s \\ &\quad + \int_0^{t \wedge \tau} \int_B c(s, Y_{s-}, x) (\mu(ds, dx) - F(dx) ds) + \int_0^{t \wedge \tau} \int_{B^c} c(s, Y_{s-}, x) \mu(ds, dx), \end{aligned}$$

showing that Y is a local strong solution to (1.4) with initial condition ℓz_0 and lifetime τ . \square

4.5. Theorem. *The following statements are true:*

- (1) *If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz and satisfy the linear growth condition, then existence and uniqueness of mild solutions to (1.3) holds.*
- (2) *If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz and locally bounded, then existence and uniqueness of local mild solutions to (1.3) holds.*
- (3) *If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz, then uniqueness of local mild solutions to (1.3) holds.*

Proof. Suppose that $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz. Let Z and Z' be two local mild solutions to (1.3) with initial conditions z_0 and z'_0 , and lifetimes τ and τ' . We define the \mathcal{H} -valued processes Y and Y' according to (4.5). By Proposition 4.4, the processes Y and Y' are local strong solutions to (1.4) with initial conditions ℓz_0 and $\ell z'_0$, and lifetimes τ and τ' , and we have $Z^\tau = \pi U Y^\tau$ and $(Z')^{\tau'} = \pi U (Y')^{\tau'}$. By Lemma 4.3, the mappings $(a, b, c|_B)$ are also locally Lipschitz, and hence, Theorem 3.12 yields that up to indistinguishability

$$Y^{\tau \wedge \tau'} \mathbb{1}_{\{\ell z_0 = \ell z'_0\}} = (Y')^{\tau \wedge \tau'} \mathbb{1}_{\{\ell z_0 = \ell z'_0\}}$$

Therefore, we have up to indistinguishability

$$Z^{\tau \wedge \tau'} \mathbb{1}_{\{z_0 = z'_0\}} = \pi U Y^{\tau \wedge \tau'} \mathbb{1}_{\{z_0 = z'_0\}} = \pi U (Y')^{\tau \wedge \tau'} \mathbb{1}_{\{z_0 = z'_0\}} = (Z')^{\tau \wedge \tau'} \mathbb{1}_{\{z_0 = z'_0\}},$$

proving uniqueness of local mild solutions to (1.3).

Now, we suppose that $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz and satisfy the linear growth condition. Let $z_0 : \Omega \rightarrow H$ be an arbitrary \mathcal{F}_0 -measurable random variable. By Lemma 4.3, the mappings $(a, b, c|_B)$ are also locally Lipschitz and satisfy the linear growth condition. Thus, by Theorem 3.14 there exists a strong solution Y to (1.4) with initial condition ℓz_0 . According to Proposition 4.4, the process $Z := \pi U Y$

is a mild solution to (1.3) with initial condition z_0 , proving the existence of mild solutions to (1.3).

If $(\alpha, \sigma, \gamma|_B)$ are locally Lipschitz and locally bounded, then a similar proof, which uses Theorem 3.15, shows that existence of local mild solutions to (1.3) holds. \square

4.6. Remark. *The structure $Z = \pi UY$ shows that mild solutions to (1.3) obtained from Theorem 4.5 have càdlàg sample paths.*

4.7. Remark. *As pointed out in [20], the existence of weak solutions to (1.3) relies on a suitable stochastic Fubini theorem. Sufficient conditions can be found in [7].*

REFERENCES

- [1] Albeverio, S., Mandrekar, V., Rüdiger, B. (2009): Existence of mild solutions for stochastic differential equations and semilinear equations with non Gaussian Lévy noise. *Stochastic Processes and Their Applications* **119**(3), 835–863.
- [2] Applebaum, D. (2005): *Lévy processes and stochastic calculus*. Cambridge University Press, Cambridge.
- [3] Cao, G., He, K., Zhang, X. (2005): Successive approximations of infinite dimensional SDEs with jump. *Stochastics and Dynamics* **5**(4), 609–619.
- [4] Da Prato, G., Zabczyk, J. (1992): *Stochastic equations in infinite dimensions*. New York: Cambridge University Press.
- [5] Davies, E. B. (1976): *Quantum theory of open systems*. London: Academic Press.
- [6] Dellacherie, C., Meyer, P. A. (1982): *Probabilités et potentiel*. Hermann: Paris.
- [7] Filipović, D., Tappe, S., Teichmann, J. (2010): Jump-diffusions in Hilbert spaces: Existence, stability and numerics. *Stochastics* **82**(5), 475–520.
- [8] Filipović, D., Tappe, S., Teichmann, J. (2010): Term structure models driven by Wiener processes and Poisson measures: Existence and positivity. *SIAM Journal on Financial Mathematics* **1**, 523–554.
- [9] Filipović, D., Tappe, S., Teichmann, J. (2012): *Invariant manifolds with boundary for jump-diffusions*. Preprint. ([arXiv:1202.1076v1](https://arxiv.org/abs/1202.1076v1))
- [10] Gawarecki, L., Mandrekar, V. (2011): *Stochastic differential equations in infinite dimensions with applications to SPDEs*. Springer, Berlin.
- [11] Gettoor, R. K. (1975): On the construction of kernels. *Séminaire de Probabilités IX*, Lecture Notes in Mathematics **465**, 443–463.
- [12] Hausenblas, E., Seidler, J. (2001): A note on maximal inequality for stochastic convolutions. *Czechoslovak Mathematical Journal* **51**(126), 785–790.
- [13] Hausenblas, E., Seidler, J. (2008): Stochastic convolutions driven by martingales: Maximal inequalities and exponential integrability. *Stochastic Analysis and Applications* **26**(1), 98–119.
- [14] Jacod, J., Shiryaev, A. N. (2003): *Limit theorems for stochastic processes*. Springer, Berlin.
- [15] Karatzas, I., Shreve, S. E. (1991): *Brownian motion and stochastic calculus*. Second Edition, Springer, New York.
- [16] Knoche, C. (2004): SPDEs in infinite dimension with Poisson noise. *Comptes Rendus Mathématique. Académie des Sciences. Paris* **339**(9), 647–652.
- [17] Kotelenez, P. (1982): A submartingale type inequality with applications to stochastic evolution equations. *Stochastics* **8**(2), 139–151.
- [18] Kotelenez, P. (1984): A stopped Doob inequality for stochastic convolution integrals and stochastic evolution equations. *Stochastic Analysis and Applications* **2**(3), 245–265.
- [19] Mandrekar, V., Rüdiger, B. (2006): Existence and uniqueness of path wise solutions for stochastic integral equations driven by Lévy noise on separable Banach spaces. *Stochastics* **78**(4), 189–212.
- [20] Marinelli, C., Prévôt, C., Röckner, M. (2010): Regular dependence on initial data for stochastic evolution equations with multiplicative Poisson noise. *Journal of Functional Analysis* **258**(2), 616–649.
- [21] Peszat, S., Zabczyk, J. (2007): *Stochastic partial differential equations with Lévy noise*. Cambridge University Press, Cambridge.
- [22] Prévôt, C. (2010): Existence, uniqueness and regularity w.r.t. the initial condition of mild solutions of SPDEs driven by Poisson noise. *Infinite Dimensional Analysis, Quantum Probability and Related Topics* **13**(1), 133–163.
- [23] Prévôt, C., Röckner, M. (2007): *A concise course on stochastic partial differential equations*. Springer, Berlin

- [24] Sz.-Nagy, B., Foiaş, C. (1970): Harmonic analysis of operators on Hilbert space. North-Holland, Amsterdam.
- [25] Taniguchi, T. (1992): Successive approximations to solutions of stochastic differential equations. *Journal of Differential Equations* **96**(1), 152–169.
- [26] Yamada, T. (1981): On the successive approximation of solutions of stochastic differential equations. *J. Math. Kyoto Univ.* **21**(3), 501–515.
- [27] Zhang, X. (2010): Stochastic Volterra equations in Banach spaces and stochastic partial differential equations. *Journal of Functional Analysis* **258**(4), 1361–1425.

LEIBNIZ UNIVERSITÄT HANNOVER, INSTITUT FÜR MATHEMATISCHE STOCHASTIK, WELFENGARTEN 1, 30167 HANNOVER, GERMANY

E-mail address: `tappe@stochastik.uni-hannover.de`